

PALEOFIRE DYNAMICS IN CENTRAL SPAIN DURING THE LATE HOLOCENE: THE ROLE OF CLIMATIC AND ANTHROPOGENIC FORCING

José Antonio López-Sáez^{1*}, Grettel Vargas², Jesús Ruiz-Fernández³, Olivier Blarquez⁴,
Francisca Alba-Sánchez², Marc Oliva⁵, Sebastián Pérez-Díaz¹,
Sandra Robles-López¹, Daniel Abel-Schaad^{1,2}

¹Archaeobiology Group, Institute of History, CCHS, CSIC, Albasanz 26-28, 28037 Madrid, Spain

²Department of Botany, University of Granada, 18071 Granada, Spain

³Department of Geography, University of Oviedo, Oviedo, Spain

⁴Département de Géographie, Université de Montréal, H3C 3J7 Montréal, Canada

⁵Institute of Geography and Spatial Planning, University of Lisbon, Lisbon, Portugal

ABSTRACT

The use of fire and, consequently, its severity and incidence on the environment, has grown steadily during the last millennia throughout the Mediterranean. This issue can be assessed in several mountain ranges of central Iberia where changes in the management policy on anthropic activities and exploitation of high-mountain environments have promoted a remarkable increase on fire frequency. Our research focuses on fire dynamics throughout the last three thousand years from three peat bog charcoal records of the Gredos range (central Iberia). Our aim is to reconstruct past fire regimes according to forest vegetation typology (*Castanea sativa*, *Pinus pinaster*, *P. sylvestris*). Charcoal influx shows low values between 3140 and 1800 cal. yr BP when forests were relatively dense both in high and mid-mountain areas. Fire appeared synchronous between 1800 and 1700 cal. yr BP for Lanzahíta and Serranillos and around 1400-1240 cal. yr BP for the three sites suggesting anthropogenic fire control between the Late Roman and the Visigothic periods that can be related to the cultivation of olive trees in the valleys and a greater human impact in high-mountain areas. By contrast, during the Muslim period (1240-850 cal. yr BP) fire dynamics becomes asynchronous. Later, fires turn again coeval in the Gredos range during the Christian period (850-500 cal. yr BP) and can be also correlated with drought phases during the Late Medieval Warm Episode. In short, our study demonstrates that fire activity has been enormously variable during the late Holocene in response to both short- and long-term regional and global climate, vegetation dynamics and land use changes.

KEY WORDS: fire history; charcoal analysis; climate variability; human impact; Gredos range; Iberian Peninsula

*Correspondence to J.A. López-Sáez, Archaeobiology Group, Institute of History, CCHS, CSIC, Albasanz 26-28, 28037 Madrid, Spain, E-mail: joseantonio.lopez@cchs.csic.es

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/ldr.2751

INTRODUCTION

Undoubtedly, the Mediterranean region is among the most impacted by land degradation (Brandt & Thornes, 1996). Its Mediterranean climate, characterized by dry summers, rough topography, weak lithology and high fire incidence have been cited as main drivers of such alterations (Vannière *et al.*, 2010; Keeley *et al.*, 2012; Marlon *et al.*, 2013; Molinari *et al.*, 2013). Nonetheless, fire has been identified as the main destabilizing element of Mediterranean terrestrial ecosystems, among human activities triggering a profound impact on this region throughout history (Le Houérou, 1973; Pausas *et al.*, 2008). In addition to natural fires, intentional burning of the forest was one of the usual management techniques employed by prehistoric societies (Naveh, 1975). In fact, it could be said that humans and fire constitute a coevolutionary system in Mediterranean high-mountain areas (López-Sáez *et al.*, 2014; Montiel & Galiana, 2016). Fire intervals are frequently <100 yr in dry biomes such as the Mediterranean region, and can even be <30 yr (Leys *et al.*, 2013). Although Mediterranean forests are able to withstand and evolve against a high fire frequency, alterations in the fire regime could have generated important consequences for past Mediterranean ecosystems (Pyne, 2009), where the role of fire has been a key factor directly affecting the distribution, structure, composition and vegetation history (Pausas, 1999; Carrión *et al.*, 2010).

High-resolution palaeoenvironmental records are required to properly understand the role of fire on the configuration and temporal dynamics of current Mediterranean landscapes, both at regional and local scales, in order to reconstruct fire regimes and predict their natural or anthropic origin (Turner *et al.*, 2008). For instance, charcoal records from peatlands or lakes have allowed to study the temporal trend of fire frequency on millennial scales, relating it to changes in vegetation and the amount of available biomass, as well as understanding its linkages to climatic variability and human impact (Conedera *et al.*, 2009; Blarquez *et al.*, 2015; Ruiz-Fernández *et al.*, 2016; Vannière *et al.*, 2016). In fact, the use of macroscopic charcoal ($\varnothing >150 \mu\text{m}$) has provided more accurate fire event reconstructions at high temporal and spatial resolution (Clark, 1988, 1989; Lynch *et al.*, 2004). Several experimental studies have shown that macrocharcoal analysis is a useful tool to determine local scale fires (500 m-1 km) since its potential transport from long distances does not alter the relationship between local fires and the sedimentary macrocharcoal record at the coring site (Clark *et al.*, 1998; Lynch *et al.*, 2004; Higuera *et al.*, 2007; Peters & Higuera, 2007).

Recent studies from the Mediterranean region have shown a clear connection between climatic shifts and millennial-scale fire regimes, particularly from the early to the mid-Holocene (11700-5000 cal. yr BP; Vannière *et al.*, 2011; Burjachs & Expósito, 2015). In contrast, during the last 3000 years, fire regime has been essentially mediated by anthropic activity (Abel-Schaad *et al.*, 2014; López-Sáez *et al.*, 2014; Morales-Molino *et al.*, 2016; Vannière *et al.*, 2016). Human impact and fire regime in the Mediterranean region have been analyzed by many research works in the last decades (Colombaroli *et al.*, 2007, 2008; Vannière *et al.*, 2008, 2011, 2016; Connor *et al.*, 2012; Leys *et al.*, 2013, 2014; Doyen *et al.*, 2015), demonstrating that fire was a key factor in mountain ecosystems management since the Neolithic with the aim of creating open spaces in forests for both agricultural and livestock activities (Colombaroli *et al.*, 2008; Lillios *et al.*, 2016). In particular, the comparative analysis of pollen and macrocharcoal records from southwestern Europe has provided substantial evidence of the role played by anthropogenic fires, which have been a major player in establishing interspecific links between plant species and shaping vegetation dynamics from mid-Holocene, causing the expansion of grasslands and scrublands, the progressive disappearance of conifer and mesophilous forests, and finally the establishment of large areas of cultivation and grazing pastures (Colombaroli *et al.*, 2008; Carrión *et al.*, 2010; Bisculm *et al.*, 2012).

Most paleofire reconstructions of the Iberian Peninsula are based on microscopic charcoal (<150 μm) from pollen slides (e.g. Burjachs & Expósito, 2015; Revelles *et al.*, 2015; Burjachs *et al.*, 2016a, 2016b; Expósito *et al.*, 2017), which mainly record a regional signal (Clark, 1988, 1989). Different methodologies, insufficient temporal resolution and, sometimes, a weak chronological control hamper the implementation of comparative studies among sites (Tinner *et al.*, 1998; Vannière *et al.*, 2016). Anyway, the incidence of human-induced fires on the frequency, fire regime and biomass burning is still an open issue in Iberia (Carrión *et al.*, 2010; Bal *et al.*, 2011; Bowman *et al.*, 2011; Morales-Molino *et al.*, 2016). Most paleofire studies have reconstructed fire history (e.g. Abel-Schaad *et al.*, 2009, 2014; López-Merino *et al.*, 2009; López-Sáez *et al.*, 2010, 2016a, 2016b; Abel-Schaad &

López-Sáez, 2013; Morales-Molino *et al.*, 2013, 2017; Burjachs & Expósito, 2015; Silva-Sánchez *et al.*, 2016) and in certain cases fire regime (e.g. Bal *et al.*, 2011; Gil-Romera *et al.*, 2014; Morales-Molino *et al.*, 2016), but only a few have taken into account regional variability on a spatial scale discerning discrepancies or agreement among records (Gil-Romera *et al.*, 2010; Vannièrè *et al.*, 2011; López-Sáez *et al.*, 2014; Morales-Molino *et al.*, 2016).

The aim of this paper is to compare three high-resolution macrocharcoal records from peat bogs located in central Spain (Gredos range, Spanish Central System). Fire variability on mountain areas regarding to anthropogenic disturbances, climate shifts and changes in vegetation cover at multi-decadal to millennial timescales can be analyzed by means of this study, performed on different sites. The data cover the late Holocene from *ca.* 3000 cal. yr BP to present. The aim is to examine the timing of fire occurrence at centennial timescales and the spatial variability associated to vegetation and location (valley bottoms vs. slopes. Five questions are addressed: i) was fire related to climate seasonality in the high-mountain Mediterranean ecosystems of the Gredos range?; ii) if so, what kind of climate control did the fire regime undergo (i.e. number of fire events within specific chronological frameworks?); iii) what was the relationship between the increase of human impact and the biomass burning pattern?; iv) can land use practices be inferred from macroscopic charcoal records?; and, v) did late Holocene fire dynamics depend on vegetation type?

MATERIAL AND METHODS

The Gredos range is found in the central part of the Iberian Central System, which divides two wide plateaus: the Northern Iberian (800-1000 m asl) and the Southern Iberian one (600-800 m asl). Its tectonic structure, a tilted horst along a south-oriented fault, establishes a distinct asymmetry between northern and southern slopes. Wet and cold winters (0°C-2°C) and dry and warm summers (20°C-22°C), as well as an average precipitation of 1400 mm per year shape climatic conditions. Regarding vegetation, holm oak (*Quercus ilex* L. subsp. *ballota* (Desf.) Sampa.) and Pyrenean oak (*Quercus pyrenaica* Willd.) spread on the mesomediterranean belt southwards, the former accompanied by xerophilous taxa like *Juniperus oxycedrus* L., *Cistus ladanifer* L., *Lavandula stoechas* Lam. and *Olea europaea* L., and the latter by mesothermophilous ones like *Arbutus unedo* L., *Corylus avellana* L., *Phillyrea angustifolia* L., *Pistacia terebinthus* L., *Lonicera implexa* Ait., *Paeonia broteroi* Boiss. & Reut. and *Viburnum tinus* L. The supramediterranean belt (> 900 m asl) is occupied by other Pyrenean oak forests. Maritime pine (*Pinus pinaster* Ait.) and sweet chestnut (*Castanea sativa* Mill.) woods also extend on the southern slopes (500-1000 m asl). Finally, the high-supra and oromediterranean belts show the wide range of grasslands and broom communities (*Cytisus oromediterraneus* Rivas-Martínez *et al.*, *Echinopartum barnadesii* (Graells) Rothm. and, to a minor extent, *Erica australis* L.) with isolated pine stands (*Pinus sylvestris* L. var. *iberica* Svoboda and *P. nigra* Arnold. subsp. *salzmannii* (Dunal) Franco) reaching the treeline. The Gredos range has historically been one of the “hot” point of forests fires in Spain (Montiel & Galiana, 2016). On one hand, there is a frequency of storms with large electrical apparatus due to the topographical characteristics of these mountains, their orientation and insolation and their situation of obstacle between the two above-mentioned plateaus; and, on the other hand, it is necessary to consider the great expansion that are having during the last fifty years the second-residence urbanizations, tourism and mountain’s sports, thanks to the forest landscape and the lush nature of the Gredos range (Galiana, 2012).

Three continuous high-resolution macrocharcoal records (16, 29 and 26 years median resolution per samples for Serranillos, Lanzahíta and Tiemblo respectively), following standardized methods of quantification, were selected in order to document the last 3000 cal. yr BP of fire regime history in the Gredos range (Figure 1). A Russian corer of 5 cm diameter was used to extract the sediment cores. The diverse sections were then sealed separately and kept on cold storage (4°C) before its handling in the laboratory. The sites were selected with the aim of discerning whether there were differences in fire history according to the type of forest. Pollen records of the three sites have already been published, in which local pollen assemblage zones were established with a constrained cluster analysis made by Coniss by transforming percentage data to its square root (Grimm, 1987). The Lanzahíta (588 m asl; mean annual precipitation 1007 mm, mean annual temperature 15.1°C; 100 cm depth) and Tiemblo (1250 m asl; mean annual precipitation 1400 mm, mean annual temperature 13°C;

120 cm depth) sequences provide the first Iberian pollen records undertaken within monospecific *Pinus pinaster* and *Castanea sativa* woodlands respectively (López-Sáez *et al.*, 2010, 2017); while that of Serranillos (1700 m asl; mean annual precipitation 937 mm, mean annual temperature 9.3°C; 120 cm depth) comes from a high-mountain area where in the last two millennia there was a *Pinus sylvestris* forest that disappeared *ca.* 500 cal. yr BP (López-Merino *et al.*, 2009; López-Sáez *et al.*, 2009). The climatic data of each record comes from Ninyerola *et al.* (2007).

Twenty one radiocarbon datings were performed on bulk peat samples (Table I). Most of these radiocarbon dates have already been published in the previously mentioned works of the authors, except one from Serranillos at 120 cm. Calib 7.1 as well as the IntCal13 curve (Reimer *et al.*, 2013) were both used to calibrate the radiocarbon dates, which were then employed by Clam 2.2 software (Blaauw, 2010) to build an age-depth model (Figure 2), with a confidence interval of 95% (2σ) through a smooth spline solution.

To reconstruct fire history, volumetric samples (1 cm^3) of sediment were collected every centimeter along all cores following the recommendations provided by Whitlock & Larsen (2001). Charcoal samples were then soaked in 15% H_2O_2 for 24 h. The sediment was subsequently sieved, collecting all charcoal particles larger than $150 \mu\text{m}$, which are assumed to have local origin ($< 10 \text{ km}$; Higuera *et al.*, 2007).

The three selected charcoal series were analyzed using the method of charcoal signal decomposition with the aim of reconstructing local fire history in each of the studied records with the CharAnalysis 1.1 software (Higuera *et al.*, 2007, 2009). These analyses were complemented with an ensemble member procedure in order to assess the sensitivity of fire event reconstruction to varying statistical methods (Blarquez *et al.*, 2013). The charcoal accumulation rate (CHAR, $\text{pieces cm}^{-2} \text{ year}^{-1}$) was established by multiplying charcoal concentration (pieces cm^{-3}) of each sample by the corresponding accretion rate (cm year^{-1}) derived from the sediment age-depth models (Figures 2 and 3). The charcoal background component (CHAR_{back}) indicates the part of the CHAR series whose temporal variation is slow in response to potential sediment mixing and even to issues arising from sampling, transport and production of charcoal (Higuera *et al.*, 2010). In order to model the CHAR_{back}, raw CHAR series (CHAR_{raw}) were filtered using the five smoothing methods available in the CharAnalysis software (Higuera *et al.*, 2009). For each method we ran 19 CHAR background simulations with variable smoothing windows equal to 150, 175, 200... 600 years. From the resulting 95 simulations, CHAR_{back} was subtracted from the CHAR_{raw} to obtain the residual high-frequency of charcoal peaks (Figure 3). This component includes the fire-induced signal (CHAR_{fire}) and the background noise not related to it but to distant and regional fires (CHAR_{noise}), as well as the noise derived from the analysis itself and from the charcoal redeposition.

A Gaussian mixture model, referred to each overlapping 500-year portions of the record, was used in each simulation to discriminate signal from noise, considered as the separation threshold the 99th percentile of the peak distribution. In this sense, CHAR_{fire} surpassing the overall variation of CHAR_{noise} is considered to be related to local fire episodes. A signal-to-noise index (SNI) was established for each sample, quantifying the discrimination between charcoal peaks related to one or multiple fire events and noise records (Figure 4). From the 95 simulations we conserved half of reconstructions (i.e. 47 reconstructions) with the higher SNI (Kelly *et al.*, 2011). From that ensemble member, we computed the distribution of the number of reconstructed event by unit of time, and determined that reconstructed fires exceeding the 75th percentile of this distribution corresponded to fire events (Figure 5) in the study area (Blarquez *et al.*, 2013).

The amount of time between detectable peaks or fire return intervals (FRIs) was calculated (Figure 6). Fire frequency was also estimated, using a Kernel density estimation technique with a 250-year bandwidth. Bootstrapped 90% confidence intervals around the fire frequency were established by bootstrap resampling of the Kernel density estimation using R “paleofire” package (Blarquez *et al.*, 2014). Changes in both FRI and fire frequency allowed us to study the diachronic variation of fire regime (Figure 6). In order to analyze the variation through time of fire disturbances, we examined FRI distributions and medians for different time intervals of the last three thousand years (Table II).

Using changes in pollen percentages as a guide, the charcoal records from Lanzahíta, Tiemblo and Serranillos were divided into zones for discussion (Table II, Figures 6 and 7). For Lanzahíta, the zones include two late Holocene *Pinus pinaster* dominated intervals between 2475-1800 and 1235-670 cal. yr BP (L1 and L3), two intervals with high values of *Cistus ladanifer* from 1800-1235 and

670-195 cal. yr BP (L2 and L4), the first one including *Olea europaea*, and a last one between 195 cal. yr BP to present (L5) with alternating values of both pollen taxa and olive tree. For Tiemblo, all zones show high percentage values of *Castanea sativa* pollen, although between 1915 and 430 cal. yr BP *Olea europaea* and anthropogenic taxa are frequent (T2-T4). Finally, for Serranillos the zones include a *Pinus sylvestris* dominated interval between 2025 and 800 cal. yr BP (S1-S4), an interval with decreasing values of pine pollen from 800-475 cal. yr BP (S5), a new *P. sylvestris* interval from 475-90 cal. yr BP (S6-S7) and an anthropozoogenic (presence of coprophilous fungi) interval from 90 cal. yr BP to present (S8).

RESULTS

Lanzahíta

CHAR averaged 0.3 pieces $\text{cm}^{-2} \text{yr}^{-1}$ between 2475 and 1800 cal. yr BP, with values ranging between 0 and 4.7 (Figure 3). High CHAR values generally prevailed between 1800 and 1235 cal. yr BP with two large peaks at 1748 and 1371 cal. yr BP (Figures 5 and 7). From 1200 cal. yr BP three periods of high CHAR are recorded with large peaks at 704, 327, 240 and -21 cal. yr BP (Figures 5 and 7). The SNI is typically > 3 (Table II, Figure 4), showing that the charcoal peak signal and the inherent noise are separated along the analyzed time series (Kelly *et al.*, 2011).

The ensemble member procedure documents 6 fire episodes (Table II, Figures 3-5). The mean FRI for the entire record is 354 years, varying amongst zones, with zone L2 = 377, L3 = 696, L4 = 218 and L5 = 261. Fire frequency shows an increasing trend from 1000 to 250 cal. yr BP where late Holocene maximum fire frequency reaching more than 4 fires by millennia (Figure 6).

Tiemblo

CHAR averaged 0.1 pieces $\text{cm}^{-2} \text{yr}^{-1}$ between 1915 and 1465 cal. yr BP, with values ranging between 0 and 1.4 (Figure 3), while between 3140-1915 cal. yr BP macrocharcoals are not documented. High CHAR values (average 0.7 pieces $\text{cm}^{-2} \text{yr}^{-1}$) are identified between 1465 and 925 cal. yr BP, with two large peaks at 1401 and 959 cal. yr BP (Figures 5 and 7). Lower CHAR reappears between 140 and 0 cal. yr BP, including a low peak at 127 cal. yr BP (Figures 5 and 7). SNI values are consistently ≥ 3 between 1915 and the present (Table II, Figure 4). However, SNI values of 0 occur between 3140-1915 cal. yr BP.

The ensemble member procedure indicates that a total of 3 fire episodes (peaks) were identified (Table II, Figures 3-5). The mean FRI interval is 637 years (Figure 6). FRIs were not determined for T1, T2, T4 and T5 zones because no fire events were recorded in those zones. Fire frequency shows maximum values between 1300 and 1100 cal. yr BP, and later a decreasing tendency until 500 cal. yr BP followed by a progressively increasing trend until the present (Figure 6).

Serranillos

CHAR averaged 0.1 pieces $\text{cm}^{-2} \text{yr}^{-1}$ between 1925 and 1775 cal. yr BP, with values ranging between 0 and 0.6 (Figure 3) including a low peak at 1852 cal. yr BP, while between 2025-1925 cal. yr BP macrocharcoals are not documented (Figures 5 and 7). Three main periods of high CHAR values are then recorded from 1450-1220, 800-475 and 100-50 cal. yr BP with large charcoal peaks at 1420, 1324, 764, 604, 492 and 108 cal. yr BP (Figures 5 and 7). The SNI is > 3 (Table II, Figure 4), showing that the charcoal peak signal and the inherent noise are separated along the analyzed time series (Kelly *et al.*, 2011).

The decomposition of the charcoal time series and associated ensemble member procedure reveals 7 fire episodes (Table II, Figures 3-5). The mean FRI for the entire record is 291 years, varying amongst zones, with zone S2 = 432, S3 = 264, S5 = 277 and S7 = 384. FRI intervals were not determined for zones S1, S4, S6 and S8 because there were too few fire events. Period with longer FRIs are documented *ca.* 1420-780 and since 500 cal. yr BP where a single event is detected at 108 cal. yr BP, whereas shorter FRIs occur from 780-490 cal. yr BP. Fire frequency progressively increase

from 1800-1400 and 1000-450 cal. yr BP and then shows a slight downward trend to the present (Figure 6).

DISCUSSION

Numerous local and regional studies have clearly demonstrated that the activity of long-term fires in the Mediterranean basin follows specific activity patterns (Colombaroli *et al.*, 2008; Turner *et al.*, 2008; Vanni re *et al.*, 2008, 2011; Gil-Romera *et al.*, 2010; Leys *et al.*, 2013; Burjachs & Exp sito, 2015). Such research provides solid evidence on synchronous trends usually interpreted as climatically-driven but also documents unforeseen ecosystem behaviors in response to particular physical configurations, vegetation and anthropic pressure. The dynamics followed by forests in the Gredos range is embedded within this sub-continental framework, demonstrating that fire activity has varied diachronically and spatially during the last two millennia (Morales-Molino *et al.*, 2013; L pez-S ez *et al.*, 2014). These facts fit current fire activity in Mediterranean ecosystems, characterized by its erratic dynamics at spatial and temporal scales depending on biomass burning, fuel availability, settlement patterns and land use strategies (Pausas *et al.*, 2008; Vanni re *et al.*, 2010, 2016; Marlon *et al.*, 2013). Although fire history of the Mediterranean basin seems to be mediated mainly by the variability of the Holocene climate (Turner *et al.*, 2008; Pyne, 2009; Vanni re *et al.*, 2011; Marlon *et al.*, 2013), several paleofire reconstructions from the Iberian Peninsula have shown that sometimes the inherent dynamics of vegetation and human activities may eclipse the climate regulating role on the fire regime, especially during the late Holocene (Gil-Romera *et al.*, 2010; Connor *et al.*, 2012; Morales-Molino *et al.*, 2013, 2016, 2017; Abel-Schaad *et al.*, 2014). For instance, in southeastern Iberian Peninsula, it has been shown that the expansion of *Pinus pinaster* forests during the mid-Holocene likewise produced significant changes in the fire regime independently of regional climate variability (Carri n & van Geel, 1999).

Considering pollen zones in the study of fire regimes in the three studied records allows us to correlate fuel types (tree species) and vegetation history with fire frequency over time. Thus, these pollen zones differ among them (Table II), so they can be interpreted as a temporal summary of average vegetation data for periods that can be considered stable within a watershed or pollen source area. This type of summary provides a large-scale examination of the control exerted by climate or by diverse disturbances of the watershed on the vegetation.

The data from Lanzah ta, Tiemblo and Serranillos peat bogs indicate that fires were infrequent prior 1800 cal. yr BP (Figures 3 and 6). Between the Iron Age and the Early Roman period the forest consisted of dense Scots pine (*Pinus sylvestris*) woodlands in the oromediterranean bioclimatic belt, as well as Maritime pine (*Pinus pinaster*) and chestnut (*Castanea sativa*) woodlands in the supramediterranean one (Figure 7). However, a low CHAR peak is observed in Serranillos at 1852 cal. yr BP (Figures 3-5), probably related to local woodland clearings by means of fire during the Early Roman period, which scarcely affected Scots pine forests. At Lanzah ta, the FRI varies during the late Holocene (Figure 6), with recorded FRIs of 261, 218, 696 and 377 years. At a regional scale, Tiemblo and Serranillos records exhibit somewhat comparable return intervals. The above-mentioned values suggest that the regional FRI was high (> 300 years) between the Iron Age and the Roman period, and progressively shorter from the Visigothic period onwards, although with remarkable differences between sites. For instance, a mean FRI of 442 years was documented from Tiemblo during the Visigothic and Muslim periods, while at Lanzah ta the FRI at this time averages 377 years (Table II). During the Christian and Early Modern periods the mean FRI is even shorter at Serranillos (277 years) and Lanzah ta (218 years), while during the Late Modern period the FRI tends to be longer at both sites. Undoubtedly, these data indicate that medieval human societies played a crucial role on the fire regime throughout the late Holocene.

Resilience of forests, human-driven and climate-influenced fires ca. 1800-1240 cal. yr BP

Increasing fire disturbance is likewise documented at Lanzahíta, Serranillos and Tiemblo *ca.* 1800-1240 cal. yr BP (Late Roman and Visigothic periods), with the rising fire frequency (Figure 6). High CHAR peaks are observed at this time in these records at 1748 and 1371 (Lanzahíta), 1401 (Tiemblo) and 1420 cal. yr BP (Serranillos). Pine decreased in Lanzahíta while olive tree (*Olea europaea*) and gum rockrose (*Cistus ladanifer*) increased from 1800 to 1235 cal. yr BP. The increase in gum rockrose may be related to more frequent fires. In fact, a sharp CHAR peak is detected at 1748 cal. yr BP (Figures 3 and 5) which coincides with maximum values of *Cistus ladanifer* and the beginning of olive cultivation in the area (Figure 7). A new high CHAR peak is documented at 1371 cal. yr BP and also corroborates a maximum percentage of gum rockrose at the end of the Visigothic period. The observed increase in fire probably benefitted *Cistus ladanifer* at Lanzahíta, given that it is sensitive to fire and is a key species helping in recolonization of *Pinus pinaster* woods after wildfires (Calvo *et al.*, 2003; Gil *et al.*, 2009). For Tiemblo, fire frequency was high up to 1000 cal. yr BP, while *Castanea sativa* values softly decrease, olive pollen appears for the first time and anthropogenic pollen taxa increases, particularly during the Visigothic period when a very sharp CHAR peak is observed at 1401 cal. yr BP (Figures 3 and 7). This generalized increase in fire disturbance can be correlated with the cultivation of olive trees at regional scale in the valleys of the Gredos range (López-Sáez *et al.*, 2014). Both olive tree growing and intensive fires could have triggered the shifts recorded on vegetation throughout zone L2, as shown by CHAR analysis of Lanzahíta sequence. The initial sharp peak of CHAR at 1748 cal. yr BP, when the first Roman settlements are reported (Martino, 2004), suggests an anthropic origin for such increase of fire regime bringing about these landscape changes in a humid phase; while the second at 1371 cal. yr BP occurs during the Visigothic period, usually considered as a point of no return for the vegetation of mountain areas of the Gredos range (López-Sáez *et al.*, 2014, 2016a).

Taking into account the above-mentioned, there is a curious fire paradox at Lanzahíta since the increase on fire frequency occurs in a humid phase during the Late Roman period (Ibero-Roman Humid Period; Figure 7; Martín-Puertas *et al.*, 2008). Additionally, fire detection from peat bogs is challenging because charcoal taphonomy in peatlands is more variable compared to lakes. Overall, only severe fire years associated to extreme droughts conditions are recorded because they can result in fires that are able to reach the coring location and burn drier peat (Conedera *et al.*, 2009; van Bellen *et al.*, 2012). If fire were related to climate factors only, such as extreme droughts, we should expect fire synchronicity at the regional scale. Here, the lack of synchronicity between the three sites rather argues that climate and extreme droughts are not the only process triggering fires and that human-induced vegetation changes and fire regime can offset the direct influence of climate. In fact, fires appear to be synchronous only *ca.* 1800-1700 cal. yr BP for Lanzahíta and Serranillos (Late Roman period) and around 1400-1300 cal. yr BP (Visigothic period) for the three sites suggesting anthropogenic fire control in relation to the cultivation of olive trees in the nearby valleys and a greater human impact in high-mountain areas. It is possible, however, that the aforementioned synchrony *ca.* 1400-1300 cal. yr BP would also be related to drier and colder climatic conditions at the beginning of the Early Medieval Cold Episode (Martín-Puertas *et al.*, 2008).

From site variability to regional fire history. Asynchronous fires ca. 1240-850 cal. yr BP

In response to cooling and dryness during the Early Medieval Cold Episode (Martín-Puertas *et al.*, 2008), pine (*Pinus sylvestris* at Serranillos and *P. pinaster* at Lanzahíta) and chestnut (Tiemblo) forests persisted between 1500 and 1000 cal. yr BP (Figure 7). Subsequently, the pine forest recovers between 1235 and 750 cal. yr BP in Lanzahíta when olive tree is no longer cultivated between the Muslim period and the beginning of the Christian period. However, gum rockrose pollen percentages remained high after fire frequency declined (Figure 7). In this sense, the increase in *Cistus ladanifer* values can also respond to the arrival of drier climatic conditions and lower temperatures related to the Early Medieval Cold Episode (1500-1000 cal. yr BP; Martín-Puertas *et al.*, 2008) that could have been beneficial to *C. ladanifer* at this site (Pérez & Moreno, 1998).

Dry climatic conditions in this chronological interval have also been documented at other sites in the Gredos range (Abel-Schaad & López-Sáez, 2013; López-Sáez *et al.*, 2014, 2016a). Meanwhile, FRIs are shorter in Lanzahíta and Tiemblo between the Visigothic and the Muslim periods (Figure 6). While in Lanzahíta and Serranillos it can be assumed that climate affected fire dynamics (fire frequency notably increases from 1000 cal. yr BP; Figure 6), although no sharp CHAR peaks are observed (Figures 3 and 5), in Tiemblo fire regime is clearly mediated by human activities since in these periods olive cultivation reaches its peak and anthropogenic pollen taxa increase their percentages (Figure 7; Morales-Molino *et al.*, 2015). In fact, a high CHAR peak is documented in Tiemblo at the end of the Muslim period (959 cal. yr BP; Figure 3). In summary, during the Muslim period (1240-850 cal. yr BP) fire frequency initially decreases and then increases (Lanzahíta, Serranillos) or remains relatively constant and then decreases (Tiemblo) (Figure 6), demonstrating that fires were asynchronous in the whole of these mountains and essentially related to human activities at Tiemblo, while in the other two records the increase on fire frequency *ca.* 1000 cal. yr BP could be correlated with the onset of the Late Medieval Warm Episode.

Grazing versus climate. Synchronous fires ca. 850-500 cal. yr BP

Fire frequency increased apparently in the Gredos range between 850 and 500 cal. yr BP as warm conditions related to the Late Medieval Warm Episode prevailed (Desprat *et al.*, 2003; Martín-Puertas *et al.*, 2008). These facts become evident in Serranillos and Lanzahíta, including several high CHAR peaks at 764, 704, 604 and 492 cal. yr BP (Figures 3 and 5) and shorter FRIs, but they do not in Tiemblo where fire frequency decreases. During the Christian period (850-500 cal. yr BP), a different landscape setting can be observed in Serranillos record with a moderate decrease of *Pinus sylvestris* forests, which coincides with increasing fire frequency and fire events (Figure 7). This is indicative of anthropogenic forest clearances by means of fire in relation to summer grazing of high-mountain areas (López-Merino *et al.*, 2009) and the Christian repopulation of the highlands of the Tiétar valley (Mariné, 1995). In contrast, the reduction of *Pinus pinaster* forests and the increase of *Cistus ladanifer* in Lanzahíta at 675 cal. yr BP do not seem to be the result of anthropic activity but rather of a dry and warmer climate. In fact, the new fire episode in Lanzahíta at 704 cal. yr BP (Figures 3 and 5) agrees with the transition between two pollen zones (L3-L4), evidencing that this episode was the driver of vegetation change. At this time, *Pinus pinaster* values drastically decrease (< 40%) while those of *Cistus ladanifer* increase at the end of the Christian period (Figure 7). The increase of gum rockrose seems to be mainly related to a warmer regional climate during the Late Medieval Warm Episode (1000-600 cal. yr BP; Desprat *et al.*, 2003; Martín-Puertas *et al.*, 2008) than with a change in the fire regime. In this sense, the observed changes in the relationship between the canopy and the understory seem to indicate some variation in the physiognomy and floristic composition of pine forests at Lanzahíta peat bog towards open woodlands due to a mixed-severity fire regime and shorter fire return intervals (696 years in L3 to 218 years in L4; Table II). In any case, burned Maritime pine forests regenerate quickly after fire (Pérez & Moreno, 1998; Calvo *et al.*, 2003), recovering high pollen values between 670 and 400 cal. yr BP (bottom of L4; Figure 7).

Human- and/or climate-mediated fires? Complexity of fire regime over the last 500 years

Fire episodes became more frequent in Lanzahíta from 400 to 195 cal. yr BP (top of L4), with two sharp CHAR peaks at 327 and 240 cal. yr BP (Figures 3 and 5) coincident with increasing *Cistus ladanifer* pollen abundance and decreasing *Pinus pinaster* values (Figure 7). Early Modern period (400-150 cal. yr BP) climatic conditions, included in the second phase of the Little Ice Age (400-100 cal. yr BP), might have heightened these effects in a period of great climatic complexity characterized by a progressive increase in temperature and extreme events of aridity and rainfall (Desprat *et al.*, 2003). These climate conditions probably caused a shortening of fire return intervals (218 years in zone L4; Table II). In contrast, no apparent high CHAR peaks are observed at this time in Serranillos and Tiemblo (Figure 3), although in the latter fire frequency shows an increasing trend from 400 cal. yr BP (Figure 6). Therefore, it is difficult to decide whether the fire regime in Lanzahíta in this period was mediated by climate or human activities, since there is no parallelism between the three macrocharcoal records.

After the fire episode at 240 cal. yr BP from Lanzahíta, a 260-year-fire-free period is documented until -21 cal. yr BP coinciding with an increase in *Pinus pinaster* values (Figure 7). In fact, during this period of decreased fire frequency from 195 cal. yr BP to the present (FRI = 261 years), this pollen type remains dominant until 100 cal. yr BP while *Cistus ladanifer* and *Olea europaea* increased from this date onwards, and a new sharp CHAR peak is observed at -21 cal. yr BP (Figure 3). The inconsistent signal displayed by *Pinus pinaster* pollen abundance during the last two hundred years could be indicating that at Lanzahíta the fire did not play a significant role in vegetation structuring during the Late Modern period, but vegetation dynamics was essentially determined by human activities.

While in Tiemblo fire frequency continues showing an increasing trend since 200 cal. yr BP, in Serranillos the opposite happens (Figure 6). These facts can be correlated with several fires occurring near Tiemblo record during the 18th and 19th centuries (Palacios, 2013). At Serranillos, a last high CHAR peak is observed at 108 cal. yr BP, more or less contemporary of another documented in Tiemblo at 127 cal. yr BP. While in Tiemblo this new fire episode hardly influenced the chestnut forest (FRI = 832 years), in Serranillos the incidence of fires and a high pastoral pressure (maximum values of coprophilous fungi; Figure 7) led to the disappearance of Scots pine forests at 500 cal. yr BP, whose percentages below 30% indicate a regional rather than a local origin (López-Sáez *et al.*, 2013). These data seem to suggest that the increase of human-mediated fires and livestock grazing in Serranillos was connected to the decline of Scots pine in the oromediterranean bioclimatic belt (López-Sáez *et al.*, 2016c). Vázquez & Moreno (2001) showed that in the Gredos range the fire cycle has been accelerated in the 20th century as a result of the dynamics introduced by fire, documenting an average fire return interval of 6.4 years in burned areas. Scots pine is considered an opportunistic species with high resilience to disturbance thanks to their light-demanding seedlings (Keeley *et al.*, 2012). However, experimental studies of the fire ecology in Iberian *Pinus sylvestris* populations have shown that this species shows significant problems to regenerate after intense fires (Escudero *et al.*, 1997, 1999; Retana *et al.*, 2002; Rodrigo *et al.*, 2004) as it does not have serotine pine cones (Tapias *et al.*, 2004), dispersing their seeds between late winter and early spring; thus, a greater frequency of forest fires in summer entails the disappearance of most seedlings limiting the regeneration of this species and affecting its distribution area (Núñez *et al.*, 2008). These experiments also demonstrate that Scots pine finds many difficulties of regeneration in high-mountain pasturelands when competing with diverse herbaceous species (Castro *et al.*, 2002).

CONCLUSIONS

The landscape of the Gredos range has behaved during the last millennia as a dynamic system thanks to its topographical heterogeneity, varied vegetation, human settlement patterns and fire history, containing different types of forests with distinct susceptibilities to fire. The sediment-charcoal analysis of three peat bog records has allowed unraveling the fire history of the Gredos range (Iberian Central System, central Spain) during the late Holocene. Understanding the complex interactions between fire, climate and anthropic activity is of paramount importance, in order to establish measures of protection against landscape degradation. Paleofire histories for the three sites demonstrate coeval changes on fire frequency, similar FRI distributions and millennial-scale synchrony. However, the fire signal also shows substantial differences among the three sites, which suggests that the fire regime also depends on the environmental gradient and local type factors separating warmer southern (Lanzahíta) from cooler sites to the north (Tiemblo), and both from the one located at higher altitude (Serranillos).

The regional susceptibility to fire of each of the studied vegetation types in the Gredos range appears to be the result of several factors, sometimes complementary, such as ignition trends, climatic variability, forest management practices and floristic composition, as well as fire-fighting strategies and effectiveness (Viedma, 2008; Moreno *et al.*, 2011). CHAR background and pollen data indicate that forests were relatively dense and capable to withstanding fires between 3140 and 1800 cal. yr BP in both the oromediterranean and the supramediterranean bioclimatic belts. Synchrony between 1800 and 1240 cal. yr BP suggests that fire regime was controlled by regional anthropic activities during

the Late Roman and Visigothic periods, particularly by olive growing in the valleys and a greater human impact in high-mountain areas. Asynchronous fire episodes between 1240 and 850 cal. yr BP suggest, instead, that local controls on fire activity, such as fuel availability and orography, were important during the Muslim period. These facts seem to rather respond to existing differences in vegetation composition, climatic conditioning, as well as to the heterogeneity of the settlement pattern and land use in each of the studied records.

One of the most noteworthy results of our paleofire analysis has been the documentation of a period in which fire episodes are synchronous between 850 and 500 cal. yr BP (Christian period), with increases in CHAR background, fire frequency and fire events, especially at Lanzahíta and Serranillos, during the warmth of the Late Medieval Warm Episode. However, such synchrony does not only respond to climatic variability but is rather due to high human impact in the case of Serranillos. Very low CHAR at Tiemblo could be related to a certain type of limitation on the spread of fire at mid elevations or reduced forest fuel availability, for example under cooler summer temperatures. It could be also linked to the fact that chestnut forest is constantly under anthropic management for the exploitation of its fruits (chestnuts) and timber, making it more difficult to burn (Seijo *et al.*, 2016). Fire has been also a major disturbance affecting Gredos range landscapes during the last three hundred years (Modern period), when both fire frequency and fire episodes have increased in these mountains. The reason could lie on the progressive land abandonment and degradation, with the consequent proliferation of shrubs, and the development of afforestation policies of former pasturelands, leading to the increase of accumulated fuel biomass.

Resilience to fire in the Gredos range appears to be mediated by the interaction between the severity and duration of each fire episode and the adaptive traits of forest species involved in the postfire response, but also by the particular human history of each area and land-use patterns. Our paleofire record suggests that the most fire-resilient forests were those that showed a rapid recovery against high fire frequency and fire events, i.e. *Pinus pinaster* in Lanzahíta, while those associated to low combustibility areas (i.e. *Castanea sativa* in Tiemblo) show a less obvious fire behavior further than the effects of fire on their forests are lower than in conifer ones. In contrast, human-driven land use changes in high-mountain environments (i.e. Serranillos) interacted with physical environment and fire in the last three hundred years, supporting the spatio-temporal dynamics of *Pinus sylvestris* forests extinction when a susceptible threshold was reached.

ACKNOWLEDGEMENTS

This research has been funded through the Desirè-HAR2013-43701-P (Plan Nacional I+D+I, Spanish Ministry of Economy and Competitiveness) and Relicflora-P11-RNM-7033 (Excellence Research Projects Program from the Andalusian Government) projects.

REFERENCES

- Aaby B, Berglund BE. 1986. Characterization of the peat and lake deposits. In: Handbook of Holocene palaeoecology and palaeohydrology, Berglund BE (Ed). Wiley, Chichester, pp. 231–246.
- Abel-Schaad D, López-Sáez JA. 2013. Vegetation changes in relation to fire history and human activities at the Peña Negra mire (Béjar Range, Iberian Central Mountain System, Spain) during the past 4.000 years. *Vegetation History and Archaeobotany* **22**: 199–214. DOI: 10.1007/s00334-012-0368-9.
- Abel-Schaad D, Hernández AM, López-Merino L, Pulido FJ, López-Sáez JA. 2009. Cabras y quemorros: Tres siglos de cambios en el paisaje de la vertiente extremeña de la Sierra de Gredos. *Revista de Estudios Extremeños* **65**: 449–478.
- Abel-Schaad D, López-Sáez JA, Pulido FJ. 2014. Heathlands, fire and grazing. A paleoenvironmental view of Las Hurdes (Cáceres, Spain) history during the last 1200 years. *Forest Systems* **23**: 247–258. DOI: 10.5424/fs/2014232-03514.
- Bal MC, Pélachs A, Pérez-Obiol R, Julià R, Cunill R. 2011. Fire history and human activities during the last 3300 cal yr BP in Spain's Central Pyrenees: The case of the Estany de Burg. *Palaeogeography, Palaeoclimatology, Palaeoecology* **300**: 179–190. DOI: 10.1016/j.palaeo.2010.12.023.
- Bisculm M, Colombaroli D, Vescovi E, Van Leeuwen JFN, Henne PD, Rothen J, Procacci G, Pasta S, La Mantia T, Tinner W. 2012. Holocene vegetation and fire dynamics in the supra-mediterranean belt of the Nebrodi Mountains (Sicily, Italy). *Journal of Quaternary Science* **27**: 687–698. DOI: 10.1002/jqs.2551.
- Blaauw M. 2010. Methods and code for classical age-modelling of radiocarbon sequences. *Quaternary Geochronology* **5**: 512–518. DOI: 10.1016/j.quageo.2010.01.002.
- Blarquez O, Girardin MP, Leys B, Ali AA, Aleman JC, Bergeron Y, Carcaillet C. 2013. Paleofire reconstruction based on an ensemble-member strategy applied to sedimentary charcoal. *Geophysical Research Letters* **40**: 2667–2672. DOI: 10.1002/grl.50504.
- Blarquez O, Vannièrè B, Marlon JR, Daniau AL, Power MJ, Brewer S, Bartlein PJ. 2014. Paleofire: an R package to analyse sedimentary charcoal records from the Global Charcoal Database to reconstruct past biomass burning. *Computers and Geosciences* **72**: 255–261. DOI: 10.1016/j.cageo.2014.07.020.
- Blarquez O, Ali AA, Girardin MP, Grondin P, Fréchette B, Bergeron Y, Hély C. 2015. Regional paleofire regimes affected by non-uniform climate, vegetation and human drivers. *Scientific Reports* **5**: 13356. DOI: 10.1038/srep13356.
- Bowman DMJS, Balch J, Artaxo P, Bond WJ, Cochrane MA, D'Antonio CM, De Fries R, Johnston FH, Keeley JE, Krawchuk MA, Kull CA, Mack M, Moritz MA, Pyne S, Ross CI, Scott AC, Sodhi NS, Swetnam TW. 2011. The human dimension of fire regime on earth. *Journal of Biogeography* **38**: 2223–2236. DOI: 10.1111/j.1365-2699.2011.02595.x.
- Brandt J, Thornes JB. 1996. Mediterranean Desertification and Land Use. Wiley, Chichester.
- Burjachs F, Expósito I. 2015. Charcoal and pollen analysis: Examples of Holocene fire dynamics in Mediterranean Iberian Peninsula. *Catena* **135**: 340–349. DOI: 10.1016/j.catena.2014.10.006.
- Burjachs F, Jones SE, Giralte S, Fernández-López de Pablo J. 2016a. Lateglacial to Early Holocene recursive aridity events in the SE Mediterranean Iberian Peninsula: The Salines playa lake case study. *Quaternary International* **403**: 187–200. DOI: 10.1016/j.quaint.2015.10.117.
- Burjachs F, Pérez-Obiol R, Picornell-Gelabert L, Revelles J, Servera-Vives G, Expósito I, Yll EI. 2016b. Overview of environmental changes and human colonization in the Balearic Islands (Western Mediterranean) and their impacts on vegetation composition during the Holocene. *Journal of Archaeological Science: Reports*. DOI: 10.1016/j.jasrep.2016.09.018.
- Calvo L, Santalla S, Marcos E, Valbuena L, Tárrega R, Luis-Calabuig E. 2003. Regeneration after wildfire in communities dominated by *Pinus pinaster*, an obligate seeder, and others dominated by *Quercus pyrenaica*, a typical resprouter. *Forest Ecology & Management* **184**: 209–223. DOI: 10.1016/S0378-1127(03)00207-X.
- Carrión JS, Fernández S, González-Sampériz P, Gil-Romera G, Badal E, Carrión-Marco Y, López-Merino L, López-Sáez JA, Fierro E, Burjachs F. 2010. Expected trends and surprises in the Lateglacial and Holocene vegetation history of the Iberian Peninsula and Balearic Islands. *Review of Palaeobotany and Palynology* **162**: 458–475. DOI: 10.1016/j.revpalbo.2009.12.007.
- Carrión JS, van Geel B. 1999. Fine-resolution upper Weichselian and Holocene palynological record from Navarrés (Valencia, Spain) and a discussion about factors of Mediterranean forest succession. *Review of Palaeobotany and Palynology* **106**: 209–236. DOI: 10.1016/S0034-6667(99)00009-3.
- Castro J, Zamora R, Hódar JA. 2002. Mechanisms blocking *Pinus sylvestris* colonization of Mediterranean mountain meadows. *Journal of Vegetation Science* **13**: 725–731. DOI: 10.1111/j.1654-1103.2002.tb02100.x.
- Clark JS. 1988. Particle motion and the theory of charcoal analysis: source area, transport, deposition and sampling. *Quaternary Research* **30**: 67–80. DOI: 10.1016/0033-5894(88)90088-9.
- Clark JS. 1989. Ecological disturbance as a renewal process: theory and application to fire history. *Oikos* **56**: 17–30. DOI: 10.2307/3566083.

- Clark, JS, Lynch J, Stocks BJ, Goldammer JG. 1998. Relationships between charcoal particles in air and sediments in west-central Siberia. *The Holocene* **8**: 19–29.
- Colombaroli, D, Marchetto A, Tinner W. 2007. Long-term interactions between Mediterranean climate, vegetation and fire regime at Lago di Massaciuccoli (Tuscany, Italy). *Journal of Ecology* **95**: 755–770. DOI: 10.1111/j.1365-2745.2007.01240.x.
- Colombaroli D, Vanni re B, Chapron E, Magny M, Tinner W. 2008. Fire-vegetation interactions during the Mesolithic-Neolithic transition at Lago dell’Accesa, Tuscany, Italy. *The Holocene* **18**: 679–692. DOI: doi: 10.1177/0959683608091779.
- Conedera M, Tinner W, Nelf C, Meurer M, Dickens AF, Krebs P. 2009. Reconstructing past fire regimes: methods, applications, and relevance to fire management and conservation. *Quaternary Science Reviews* **28**: 555–576. DOI: 10.1016/j.quascirev.2008.11.005.
- Connor SE, Ara jo J, van der Knapp WO, van Leeuwen JFN. 2012. A long-term perspective on biomass burning in the Serra da Estrela, Portugal. *Quaternary Science Reviews* **55**: 114–124. DOI: 10.1016/j.quascirev.2012.08.007.
- Desprat S, S nchez-Go ni MF, Loutre MF. 2003. Revealing climatic variability of the last three millennia in northwestern Iberia using pollen influx data. *Earth and Planetary Science Letters* **213**: 63–78. DOI: 10.1016/S0012-821X(03)00292-9.
- Doyen E, Vanni re B, Rius D, B geot C, Millet L. 2015. Climate and biomass control on fire activity during the late-glacial/early Holocene transition in temperate ecosystems of the upper Rhone valley (France). *Quaternary Research* **83**: 94–104. DOI: 10.1016/j.yqres.2014.08.004.
- Escudero A, Barrero S, Pita JM. 1997. Effects of high temperatures and ash on seed germination of two Iberian pines (*Pinus nigra* ssp *salzmannii*, *P. sylvestris* var *iberica*). *Annales des Sciences Foresti res* **54**: 553–561. DOI: 10.1051/forest:19970605.
- Escudero A, Sanz MV, Pita JM, P rez-Garc a F. 1999. Probability of germination after heat treatment of native Spanish pines. *Annals of Forest Science* **56**: 511–520. DOI: 10.1051/forest:19990608.
- Exp sito I, Burjachs F, Allu , E. 2017. Filling in the gaps: The contribution of non-pollen palynomorphs to knowledge about the local environment of the Sierra de Atapuerca caves during the Pleistocene. *Quaternary International* **433A**: 221–242. DOI: 10.1016/j.quaint.2015.09.016.
- Galiana L. 2012. Las interfaces urbano-forestales: un nuevo territorio de riesgo en Espa a. *Bolet n de la Asociaci n de Ge grafos Espa oles* **58**: 205–226.
- Gil L, L pez R, Garc a-Mateos A, Gonz lez-Doncel I. 2009. Seed provenance and fire-related reproductive traits of *Pinus pinaster* in central Spain. *International Journal of Wildland Fire* **18**: 1003–1009. DOI: 10.1071/WF08101.
- Gil-Romera G, Carri n JS, Pausas JG, Sevilla M, Lamb HJ, Fern ndez S, Burjachs F. 2010. Holocene fire activity and vegetation response in South-Eastern Iberia. *Quaternary Science Reviews* **29**: 1082–1092. DOI: 10.1016/j.quascirev.2010.01.006.
- Gil-Romera G, Gonz lez-Samp riz P, Lasheras- lvarez L, Sevilla-Callejo M, Moreno A, Valero-Garc s B, L pez-Merino L, Carri n JS, P rez Sanz A, Aranbarri J, Garc a-Prieto E. 2014. Biomass-modulated fire dynamics during the Last Glacial-Interglacial transition at the Central Pyrenees (Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology* **402**: 113–124. DOI: 10.1016/j.palaeo.2014.03.015.
- Grimm EC. 1987. Coniss: a Fortran 77 program for stratigraphically constrained cluster analysis by the method of incremental sum of squares. *Computers & Geosciences* **13**: 13–35. DOI: 10.1016/0098-3004(87)90022-7.
- Higuera PE, Peters ME, Brubaker LB, Gavin DG. 2007. Understanding the origin and analysis of sediment-charcoal records with a simulation model. *Quaternary Science Reviews* **26**: 1790–1809. DOI: 10.1016/j.quascirev.2007.03.010.
- Higuera PE, Brubaker LB, Anderson PM, Hu FS, Brown TA. 2009. Vegetation mediated the impacts of postglacial climate change on fire regimes in the south-central Brooks Range, Alaska. *Ecological Monographs* **79**: 201–219. DOI: 10.1890/07-2019.1.
- Higuera PE, Gavin DG, Bartlein PJ, Hallett DJ. 2010. Peak detection in sediment-charcoal records: impacts of alternative data analysis methods on fire-history interpretations. *International Journal of Wildland Fire* **19**: 996–1014. DOI: 10.1071/WF09134.
- Keeley JE, Bond W, Bradstock R, Pausas J, Rundel P. 2012. *Fire in Mediterranean Ecosystems*. Cambridge University Press, London.
- Kelly RF, Higuera PE, Barrett CM, Hu FS. 2011. A signal-to-noise index to quantify the potential for peak detection in sediment-charcoal records. *Quaternary Research* **75**: 11–17. DOI: 10.1016/j.yqres.2010.07.011.
- Le Hou rou NH. 1973. Fire and vegetation in the Mediterranean basin. In: *Proceedings of the 13th Annual Tall Timber Fire Ecology Conference*. Tall Timber Research Station, Tallahassee, Florida, pp. 237–277.
- Leys B, Carcaillet C, Dezileau L, Ali AA, Bradshaw RHW. 2013. A comparison of charcoal measurements for reconstruction of Mediterranean paleo-fire frequency in the mountains of Corsica. *Quaternary Research* **79**: 337–349. DOI:10.106/j.yqres.2013.01.003.

- Leys B, Finsinger W, Carcaillet C. 2014. Historical range of fire frequency is not the Achilles' heel of the Corsican black pine ecosystem. *Journal of Ecology* **102**: 381–395. DOI: 10.1111/1365-2745.12207.
- Lillios KT, Blanco-González A, Lee B, López-Sáez JA. 2016. Mid-Late Holocene climate, demography, and cultural dynamics in Iberia: a multi-proxy approach. *Quaternary Science Reviews* **135**: 138–153. DOI: 10.1016/j.quascirev.2016.01.011.
- López-Merino L, López-Sáez JA, Alba-Sánchez F, Pérez-Díaz S, Carrión JS. 2009. 2000 years of pastoralism and fire shaping high-altitude vegetation of Sierra de Gredos in central Spain. *Review of Palaeobotany and Palynology* **158**: 42–51. DOI: 10.1016/j.revpalbo.2009.07.003.
- López-Sáez JA, López-Merino L, Alba-Sánchez F, Pérez-Díaz S. 2009. Contribución paleoambiental al estudio de la trashumancia en el sector abulense de la Sierra de Gredos. *Hispania. Revista Española de Historia* **231**: 9–38. DOI: 10.3989/hispania.2009.v69.i231.97.
- López-Sáez JA, López-Merino L, Alba-Sánchez F, Pérez-Díaz S, Abel-Schaad D, Carrión JS. 2010. Late Holocene ecological history of *Pinus pinaster* forests in the Sierra de Gredos of central Spain. *Plant Ecology* **206**: 195–209. DOI: 10.1007/s11258-009-9634-z.
- López-Sáez JA, Sánchez-Mata D, Alba-Sánchez F, Abel-Schaad D, Gavilán RG, Pérez-Díaz S. 2013. Discrimination of Scots pine forests in the Iberian Central System (*Pinus sylvestris* var. *iberica*) by means of pollen analysis. Phytosociological considerations. *Lazaroa* **34**: 191–208. DOI: 10.5209/rev_LAZA.2013.v34.n1.43599.
- López-Sáez JA, Abel-Schaad D, Pérez-Díaz S, Blanco-González A, Alba-Sánchez F, Dorado M, Ruiz-Zapata B, Gil-García MJ, Gómez C, Franco-Múgica F. 2014. Vegetation history, climate and human impact in the Spanish Central System over the last 9,000 years. *Quaternary International* **353**: 98–122. DOI: 10.1016/j.quaint.2013.06.034.
- López-Sáez JA, Abel-Schaad D, Robles-López S, Pérez-Díaz S, Alba-Sánchez F, Nieto-Lugilde D. 2016a. Landscape dynamics and human impact on high-mountain woodlands in the western Spanish Central System during the last three millennia. *Journal of Archaeological Science: Reports* **9**: 203–218. DOI: 10.1016/j.jasrep.2016.07.027.
- López-Sáez JA, Alba-Sánchez F, Robles-López S, Pérez-Díaz S, Abel-Schaad D, Sabariego S, Glais A. 2016b. Exploring seven hundred years of transhumance, climate dynamic, fire and human activity through a historical mountain pass in central Spain. *Journal of Mountain Science* **13**: 1139–1153. DOI: 1007/s11629-010-3885-7.
- López-Sáez JA, Sánchez-Mata D, Gavilán RG. 2016c. Syntaxonomical update on the relict groves of Scots pine (*Pinus sylvestris* L. var. *iberica* Svoboda) and Spanish black pine (*Pinus nigra* Arnold subsp. *salzmannii* (Dunal) Franco) in the Gredos range (central Spain). *Lazaroa* **37**: 153–172. DOI: 10.5209/LAZA.54043.
- López-Sáez JA, Glais A, Robles-López S, Alba-Sánchez F, Pérez-Díaz S, Abel-Schaad D, Luelmo-Lautenschlaeger R. 2017. Unraveling the naturalness of sweet chestnut forests (*Castanea sativa* Mill.) in central Spain. *Vegetation History and Archaeobotany*. DOI: 10.1007/s00334-016-0575-x.
- Lynch JA, Clark JS, Stocks BJ. 2004. Charcoal production, dispersal, and deposition from the Fort Providence experimental fire: Interpreting fire regimes from charcoal records in boreal forests. *Canadian Journal of Forest Research* **34**: 1642–1656. DOI: 10.1139/x04-071.
- Mariné M. 1995. El patrimonio arqueológico de la Sierra de Gredos. In: Gredos: territorio, sociedad y cultura, Trotitiño MA (Ed.). Diputación de Ávila, Ávila, pp. 19–48.
- Marlon JR, Bartlein PJ, Daniau AL, Harrison SP, Maezumi SY, Power MJ, Tinner W, Vannière B. 2013. Global biomass burning: a synthesis and review of Holocene paleofire records and their controls. *Quaternary Science Reviews* **65**: 5–25. DOI: 10.1016/j.quascirev.2012.11.029.
- Martín-Puertas C, Valero-Garcés BL, Mata MP, González-Sampérez P, Bao R, Moreno A, Stefanova V. 2008. Arid and humid phases in southern Spain during the last 4000 years: The Zoñar Lake record, Córdoba. *The Holocene* **18**: 907–921. DOI: 10.1177/0959683608093533.
- Martino D. 2004. Tierras con historia: Lanzahíta. In: Lanzahíta (Ávila). Historia, naturaleza, tradiciones, González JM, Chavarría JA, López-Sáez JA (Eds.). SEVAT, Madrid, pp. 41–74.
- Molinari C, Lehsten V, Bradshaw RHW, Power MJ, Harmand P, Arneith A, Kaplan JO, Vannière B., Sykes MT. 2013. Exploring potential drivers of European biomass burning over the Holocene: a data-model analysis. *Global Ecology and Biogeography* **22**: 1248–1260. DOI: 10.1111/geb.12090.
- Montiel C, Galiana L. 2016. Fire scenarios in Spain: a territorial approach to proactive fire management in the context of global change. *Forests* **7**: 273. DOI: 10.3390/f7110273.
- Morales-Molino C, García-Antón M, Postigo-Mijarra JM, Morla C. 2013. Holocene vegetation, fire and climate interactions on the westernmost fringe of the Mediterranean Basin. *Quaternary Science Reviews* **59**: 5–17. DOI: 10.1016/j.quascirev.2012.10.027.
- Morales-Molino C, Vescovi E, Krebs P, Carlevaro E, Kaltenrieder P, Conedera M, Colombaroli D. 2015. The role of human-induced fire and sweet chestnut (*Castanea sativa* Mill.) cultivation on the long-term

- landscape dynamics of the southern Swiss Alps. *The Holocene* **25**: 482–494. DOI: 10.1177/0959683614561884.
- Morales-Molino C, Tinner W, García-Antón M, Colombaroli D. 2016. The historical demise of *Pinus nigra* forests in the Northern Iberian Plateau (south-western Europe). *Journal of Ecology*. DOI: 10.1111/1365-2745.12702.
- Morales-Molino C, Colombaroli D, Valbuena-Carabaña M, Tinner W, Salomón RL, Carrión JS, Gil L. 2017. Land-use history as a major driver for long-term forest dynamics in the Sierra de Guadarrama National Park (central Spain) during the last millennia: implications for forest conservation and management. *Global and Planetary Change* **152**: 64–75. DOI: 10.1016/j.gloplacha.2017.02.012.
- Moreno JM, Viedma O, Zavala G, Luna B. 2011. Landscape variables influencing forest fires in central Spain. *International Journal of Wildland Fire* **20**: 678–689. DOI: 10.1071/WF10005.
- Naveh Z. 1975. The evolutionary significance of fire in the Mediterranean region. *Vegetatio* **29**: 199–208. DOI: 10.1007/BF02390011.
- Ninyerola M, Pons X, Roure M. 2007. Atlas climatic digital de la Península Ibérica. Metodología y aplicaciones en bioclimatología y geobotánica. Universidad Autónoma de Barcelona, Bellaterra.
- Núñez MR, Calvo L, Pando V, Bravo F. 2008. Floristic changes induced by fire on *Pinus sylvestris* plantations in northwestern of Spain. *Investigación Agraria: Sistemas y Recursos Forestales* **17**: 168–177.
- Palacios MT. 2013. Fuentes documentales para el estudio de los fuegos forestales en Ávila. In: Presencia histórica del fuego en el territorio, Montiel-Molina C (Ed.). M.I.M.A.M., Madrid, pp. 155–175.
- Pausas JC. 1999. Mediterranean vegetation dynamics: modelling problems and functional types. *Plant Ecology* **140**: 27–39. DOI: 10.1023/A:1009752403216.
- Pausas JC, Llovet J, Rodrigo A, Vallejo R. 2008. Are wildfires a disaster in the Mediterranean basin? A review. *International Journal of Wildland Fire* **17**: 713–723. DOI: 10.1071/WF07151.
- Pérez B, Moreno JM. 1998. Fire-type and forestry management effects on the early postfire vegetation dynamics of a *Pinus pinaster* woodland. *Plant Ecology* **134**: 27–41. DOI: 10.1023/A:1009733818670.
- Peters ME, Higuera PE. 2007. Quantifying the source area of macroscopic charcoal with a particle dispersal model. *Quaternary Research* **67**: 304–310. DOI: 10.1016/j.yqres.2006.10.004.
- Pyne SJ. 2009. Eternal flame: an introduction to the fire history of the Mediterranean. In: Earth Observation of Wildland Fires in Mediterranean Ecosystems, Chuvieco E (Ed.). Springer-Verlag, Berlin-Heidelberg, pp. 11–26.
- Reimer PJ, Bard E, Bayliss A, Beck JW, Blackwell PG, Bronk Ramsey C, Buck CE, Cheng H, Edwards RL, Friedrich M, Grootes PM, Guilderson TP, Haflidason H, Hajdas I, Hatté C, Heaton TJ, Hoffmann DL, Hogg AG, Hughen KA, Kaiser KF, Kromer B, Manning SW, Niu M, Reimer RW, Richards DA, Scott EM, Southon JR, Staff RA, Turney CSM, van der Plicht J. 2013. Intcal13 and marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* **55**: 1869–1887. DOI: 10.2458/azu_js_rc.55.16947.
- Retana J, Espelta JM, Habrouk A, Ordóñez JL, Solà-Morales F. 2002. Regeneration patterns of three Mediterranean pines and forest changes after a large wild-fire in northeastern Spain. *Ecoscience* **9**: 89–97. DOI: 10.1080/11956860.2002.11682694.
- Revelles J, Cho S, Iriarte E, Burjachs F, van Geel B, Palomo A, Piqué R, Peña-Chocarro L, Terradas X. (2015). Mid-Holocene vegetation history and Neolithic land-use in the Lake Banyoles area (Girona, Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology* **435**: 70–85. DOI: 10.1016/j.palaeo.2015.06.002.
- Rodrigo A, Retana J, Xavier F. 2004. Direct regeneration is not the only response of Mediterranean forests to large fires. *Ecology* **85**: 716–729. DOI: 10.1890/02-0492.
- Ruiz-Fernández J, Nieuwendam A, Oliva M, Lopes V, Cruces A, Janeiro A, Freitas C, López-Sáez JA. 2016. Cryogenic processes and fire activity in a high Atlantic mountain area in NW Iberia (Picos de Europa) during the Mid-Late Holocene. *Science of the Total Environment* **573**: 1159–1170. DOI: 10.1016/j.scitotenv.2016.03.022.
- Seijo F, Millington JDA, Gray R, Hernández L, Sangüesa-Barreda G, Camarero JJ. 2016. Divergent fire regimes in two contrasting Mediterranean chestnut forest landscapes. *Human Ecology*. DOI: 10.1007/s10745-016-9879-9.
- Silva-Sánchez N, Martínez-Cortizas A, Abel-Schaad D, López-Sáez JA, Mighall TM. 2016. Influence of climate change and human activities in the organic and inorganic composition of peat during the Little Ice Age (El Payo mire, Gata range, W Spain). *The Holocene* **26**: 1290–1303. DOI: 10.1177/0959683616638439.
- Tapias R, Climent J, Pardos J, Gil L. 2004. Life histories of Mediterranean pines. *Plant Ecology* **171**: 53–68. DOI: 10.1023/B:VEGE.0000029383.72609.f0.
- Tinner W, Conedera M, Ammann B, Gaggeler HW, Gedye S, Jones R, Sagesser B. 1998. Pollen and charcoal in lake sediments compared with historically documented forest fires in southern Switzerland since AD 1920. *The Holocene* **8**: 31–42. DOI: 10.1191/095968398667205430.
- Turner R, Roberts N, Jones MD. 2008. Climatic pacing of Mediterranean fire histories from lake sedimentary microcharcoal. *Global and Planetary Change* **63**: 317–324. DOI: 10.1016/j.gloplacha.2008.07.002.

- van Bellen S, Garneau M, Ali AA, Bergeron Y. 2012. Did fires drive Holocene carbon sequestration in boreal ombrotrophic peatlands of eastern Canada? *Quaternary Research* **78**: 50–59. DOI: 10.1016/j.yqres.2012.03.009.
- Vanni re B, Colombaroli D, Chapron E, Leroux A, Tinner W, Magny M. 2008. Climate versus human-driven fire regimes in Mediterranean landscapes: the Holocene record of Lago dell'Accesa (Tuscany, Italy). *Quaternary Science Reviews* **27**: 1181–1196. DOI: 10.1016/j.quascirev.2008.02.011.
- Vanni re B, Colombaroli D, Roberts N. 2010. A fire paradox around the Mediterranean. *PAGES Newsletter* **18**: 63–65.
- Vanni re B, Power MJ, Roberts N, Tinner W, Carri n J, Magny M, Bartlein P, Colombaroli D, Daniau AL, Finsinger W, Gil-Romera G, Kaltenrieder P, Magri D, Pini R, Sadori L, Turner R, Valsechi V, Vescovi E. 2011. Circum-Mediterranean fire activity and climate changes during the mid-Holocene environmental transition (8500-2500 cal. BP). *The Holocene* **21**: 53–73. DOI: 10.1177/0959683610384164.
- Vanni re B, Blarquez O, Rius D, Doyen E, Br ucher T, Colombaroli D, Connor S, Feurdean A, Hickler T, Kaltenrieder P, Lemmen C, Leys B, Massa C, Olofsson J. 2016. 7000-year human legacy of elevation-dependent European fire regimes. *Quaternary Science Reviews* **132**: 206–212. DOI: 10.1016/j.quascirev.2015.11.012.
- V zquez A, Moreno JM. 2001. Spatial distribution of forest fires in Sierra de Gredos (Central Spain). *Forest Ecology & Management* **147**: 55–65. DOI: 10.1016/S0378-1127(00)00436-9.
- Viedma O. 2008. The influence of topography and fire in controlling landscape composition and structure in Sierra de Gredos (Central Spain). *Landscape Ecology* **23**: 657–672. DOI: 10.1007/s10980-008-9228-5.
- Whitlock C, Larsen C. 2001. Charcoal as a fire proxy. In: Tracking environmental change using lake sediments, vol. 3, Terrestrial, algal, and siliceous indicators, Smol JP, Birks HJB, Last WM (Eds.). Kluwer, Dordrecht, pp. 75–97.

Accepted Article

Table I. AMS-radiocarbon data with 2σ range of calibration.

Laboratory code	Depth (cm)	AMS ^{14}C age BP	Age cal yr BP (2σ)	Mean Age cal yr BP
<i>Lanzahíta</i>				
CNA-095	7.5	Modern	-	-
Ua-24879	24.5	780 ± 35	759-667	705
Ua-24878	35	1020 ± 35	1048-800	940
CNA-105	44.5	1315 ± 29	1295-1183	1260
CSIC-1877	64.5	1907 ± 35	1928-1737	1853
Ua-19515	75.5	2280 ± 55	2420-2140	2260
CSIC-1876	100	2387 ± 32	2677-2345	2413
<i>Serranillos</i>				
CSIC-1767	39.5	522 ± 27	624-509	535
CSIC-1768	82.5	1664 ± 35	1693-1420	1569
Ua-21490	91	1895 ± 45	1929-1717	1839
CSIC-1938	102.5	1938 ± 35	1986-1819	1888
Poz-83925	120	2050 ± 30	2113-1932	2013
<i>Tiemblo</i>				
CNA-211	7	210 ± 60	429-0	192
CNA-185	15	375 ± 40	506-316	432
CNA-103	29	815 ± 35	786-680	725
CNA-186	41	1130 ± 35	1173-961	1031
CNA-546	55	1425 ± 35	1380-1289	1327
CNA-551	72	1875 ± 25	1876-1733	1825
CNA-540	88	2290 ± 25	2352-2184	2330
CNA-097	105	2675 ± 25	2844-2750	2775
CNA-096	120	2970 ± 30	3229-3007	3136

Accepted

Table II. Zone, mean values for selected pollen and non-pollen palynomorph percentages and CHAR and measures of fire disturbance for Lanzahíta (L1-L5), Tiemblo (T1-T6) and Serranillos (S1-S8), where cal. yr BP = calendar years before present; CHAR = charcoal accumulation rate; SNI = signal to noise index; FRI = fire return interval; N/A = not available.

Zone	Age (cal. yr BP)	CHAR (x10 ⁻¹)	SNI	Fires (no.)	FRI (years)	Pollen and non-pollen palynomorph percentages							
						<i>Pinus pinaster</i>	<i>Pinus sylvestris</i>	Genisteeae	Poaceae	<i>Castanea</i>	Coprophilous fungi	<i>Olea</i>	<i>Cistus ladanifer</i>
L5	0-195	13.51	~ 3	1	261	32-50	< 3	-	7-18	< 3	2-25	1-6	3-10
L4	195-670	10.67	≥ 3	2	218	29-65	< 3	-	4-13	< 3	2-23	-	4-16
L3	670-1235	7.79	≥ 3	1	696	53-69	< 3	-	4-7	< 3	< 3	-	2-8
L2	1235-1800	16.03	≥ 3	2	377	29-45	< 3	-	9-16	< 3	2-15	4-11	2-10
L1	1800-2475	2.95	≥ 3	0	N/A	52-62	< 3	-	5-10	< 3	< 3	-	< 3
T6	0-140	0.52	≥ 3	1	832	3-7	3-6.5	< 3	< 3	46-48.6	-	< 3	< 3
T5	140-430	0.00	≥ 3	0	N/A	< 3	< 3	< 3	< 3	37-46	5-13	< 3	< 3
T4	430-925	1.64	≥ 3	0	N/A	< 3	< 5	< 3	< 3	34-44.5	6-10	< 3	< 3
T3	925-1465	7.33	≥ 3	2	442	2-4	3-6	< 3	< 3	30.5-38	3-7	2-4	< 3
T2	1465-1915	1.08	≥ 3	0	N/A	< 2	8-10	< 3	< 3	34-41	5	0-1	< 3
T1	1915-3140	0.00	0	0	N/A	< 5	4-11	< 3	< 3	31-42	< 1	-	< 3
S8	0-90	28.22	≥ 3	0	N/A	< 1	10-20	< 3	25-50	-	5-13	5-12	< 1
S7	90-230	8.75	≥ 3	1	384	< 3	35-58	< 5	20	-	4-22	2-4	< 1
S6	230-475	11.25	≥ 3	0	N/A	1-11	20-50	-	19-49	-	< 3	1-6	< 1
S5	475-800	42.26	≥ 3	3	277	-	25-42	-	25-50	-	< 1	-	1
S4	800-1220	2.18	≥ 3	0	N/A	< 3	56-70	< 1	4-15	-	< 1	-	< 1
S3	1220-1775	7.31	≥ 3	2	264	< 3	45-67	-	5-17	-	< 1	< 1	< 1
S2	1775-1925	1.32	≥ 3	1	432	5-7	57-69	-	8-19	-	< 1	< 1	-
S1	1925-2025	0.00	≥ 3	0	N/A	5-7	60-70	< 3	< 5	-	< 1	-	-




Figure 1. Location of the studied sites in the Gredos range (central Iberia).

Accepted




Figure 2. Lithology (following Aaby & Berglund, 1986), sedimentation rate (mm/yr) and age-depth models based on calibrated ¹⁴C dates of the Lanzahíta, Tiemblo and Serranillos cores which were constructed using the Clam 2.2 software package. The black line of the plots show the best fit age-depth model and the grey envelope of the plots show the final 95% confidence.

Accepted




Figure 3. CHAR_{raw} (grey bars), interpolated CHAR (stair-step, black line) and identified fire events (+ symbol) for (A) Lanzahíta, (C) Tiemblo and (E) Serranillos.




Figure 4. Mean Signal to Noise Index (SNI) for the ensemble members (47 reconstructions) and identified fire events (+ symbol) for (A) Lanzahíta, (C) Tiemblo and (E) Serranillos.




Figure 5. Cumulative fires through time. Black triangles = Serranillos; black circles = Tiemblo and white circles = Lanzahíta.




Figure 6. Late Holocene fire disturbance history for Lanzahíta (A-B), Tiemblo (C-D) and Serranillos (E-F). Fire frequency from the kernel density estimation procedure is represented using a black line along with its bootstrapped 90 % confidence interval (grey area, A-C-E). Fire Return Intervals (FRIs) are represented using black squares (B-D-F). Also shown are zone boundaries for Lanzahíta (L2-L5), Tiemblo (T2-T6) and Serranillos (S2-S8).




Figure 7. Fire events and pollen abundances from Serranillos, Lanzahíta and Tiemblo records plotted against age (cal. yr BP) and cultural periods. Color shading correspond to climatic events discussed in the text (LIA = Little Ice Age; LMWE = Late Medieval Warm Episode; EMCE = Early Medieval Cold Episode; IRHP = Ibero-Roman Humid Period; 2.8 kyr = 2.8 cal. kyr BP event). Coprophilous fungi include *Sordaria* and *Sporormiella*. Anthropogenic taxa types include *Aster*, *Cardueae* and *Cichorioideae*. Pollen and non-pollen palynomorph values are relative to total terrestrial pollen abundance.

Accepted